



## Butyric acid fermentation with in-situ products separation

Baroi, George Nabin; Skiadas, Ioannis; Westermann, Peter; Gavala, Hariklia N.

*Publication date:*  
2014

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Baroi, G. N., Skiadas, I., Westermann, P., & Gavala, H. N. (2014). *Butyric acid fermentation with in-situ products separation*. Poster presented at Tomorrow's biorefineries in Europe, Brussels, Belgium.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



Baroi G.N<sup>1</sup>, Skiadas I.V.<sup>1</sup>, Westermann P.<sup>1</sup>, Gavala H.N<sup>1</sup>.

<sup>1</sup>Aalborg University Copenhagen (AAU-Cph), Department of Biotechnology, Chemistry and Environmental Engineering, A C Meyers Vænge 15, DK 2450 Copenhagen SV Denmark.


## Background

Butyric acid and its esters are used as food supplements, as artificial flavours, as solvent and even its potential beneficial effects in intestinal and extra intestinal diseases and sickle cell disease have been reported. Another potential use of butyric acid is to produce butanol via catalytic hydrogenation. The demand of the butyric acid is approximately 50,000 ton/year.

The main **bottlenecks of commercialization** of biological production of acids (in overall) are:


- (1) utilization of cheap feed stocks
- (2) Proper strain selection &/or improvement
- (3) Process development for higher productivity
- (4) Downstream processing

The first three aspects were successfully considered for biological butyric acid production from pretreated and hydrolysed wheat straw (PHWS) by *C.tyrobutyricum*. Continuous fermentation and in-situ separation of butyric acid increased C6 and C5 consumption and butyric acid production rates compared to batch fermentation without in-situ products removal. The process efficiency was verified in pilot-scale as well. However, further improvement of downstream process required.




Process model of butyric acid production from wheat straw


## Rates of synthetic medium (mixed glucose and xylose) based fermentation with and without in-situ acids separation by REED at HRT 1 day




## Schematic diagram of the experimental setup



## Rates of PHWS based fermentation with and without in-situ acids separation by REED at HRT 1 day



## Continuous fermentation of 100% PHWS with REED



## Results

## Conclusion

- Fermentation coupled with REED in-situ separation of PHWS with the adapted *C. tyrobutyricum* gave a higher (28 to 58%) butyric acid yield compared to synthetic medium fermentation.
- Continuous fermentation exhibited much higher (>600% for 60% PHWS) sugars consumption rates compared to batch fermentations
- With REED in-situ acid separation results higher sugars consumption rate and C4 production rates (> 46%)
- Fermentation coupled with REED in-situ separation of **100% PHWS** continued unhindered with just urea and K<sub>2</sub>HPO<sub>4</sub> added reaching a productivity of **1,31 g/l/h** butyric acid production and **97%** sugars utilization

## Acknowledgement

The authors wish to thank the Commission of the European Communities for the financial support of this work under SUPRABIO project (FP7-cooperationproject no 241640).

